



# **Mohave County Overdose Fatality Review**

## **Annual Report 2023**

Mohave County Department of Public Health  
Overdose Data to Action Program



*This publication was made possible by grant number H79TI087838 from SAMHSA. The views, opinions, and content of this publication are those of the author and do not necessarily reflect the views, opinions, or policies of SAMHSA or HHS.*

## Acknowledgements

The Overdose Fatality Review (OFR) team is comprised of key subject matter experts from a variety of local organizations. This report would not be possible without the work and dedication of its team members. A very special thank you to the following agencies:

- Kingman Regional Medical Center
- Mohave County Sheriff's Office
- Mohave County Jail
- Mohave Substance Treatment Education & Prevention Partnership (MSTEPP)
- Mohave County Medical Examiner's Office
- Bullhead City Police Department
- Lake Havasu City Police Department
- Kingman Police Department
- Mohave County Department of Public Health
- Arizona Department of Public Safety
- Mohave Mental Health
- Terros Health
- Southwest Behavioral and Health Services
- Tribal Action Planning Committee

## Prepared by

Celina Lagpacan, Program Coordinator  
Mohave County Department of Public Health  
Overdose Data to Action Program  
Publication Date: 07/01/2025



## Table of Contents


Introduction.....	4
Mission.....	5
Methodology.....	5
Data Sources.....	5
Data Overview.....	6
Demographics.....	6
Circumstances.....	7
Contributing Factors and Commonalities.....	9
Recommendations & Action Steps.....	13
Conclusion.....	15

## Introduction

During the past decade, the number and rate of drug overdose deaths in Mohave County has risen annually. In 2017, Governor Doug Ducey declared a public health state of emergency in response to the opioid epidemic leading to a heightened approach to addressing opioid misuse and abuse throughout the state. This led to the formation of locally based Overdose Fatality Review (OFR) teams in all counties across Arizona.

### 12 Month-ending Provisional Number and Percent Change of Drug Overdose Deaths

Based on data available for analysis on: May 4, 2025



- Drug overdose deaths involving psychostimulants such as methamphetamine are increasing with and without opioid involvement.
- Drug overdoses impact families, communities, workplaces, and the economy.
- **The drug overdose mortality rate per 100,000 in Arizona for 2023 is 26.3.**

[www.cdc.gov](http://www.cdc.gov)

## Mission

The mission of the Mohave County Overdose Fatality Review (OFR) Team is to conduct case reviews and gather data to report key trends found in local drug overdose fatalities. The team uses local professionals throughout the county to conduct a detailed review of the circumstances surrounding overdose fatalities. The major purpose of the program is to develop and implement data-driven recommendations for reducing preventable drug overdose deaths.

## Methodology

Arizona Department of Health Services sent a list of 60 decedents who had drug overdose (prescription or illicit) listed as their primary cause of death in 2023 to Mohave County Department of Public Health. The Mohave County OFR team selected a sample size of 27 decedents whose cause of death was attributed to prescription or illicit drugs and who had either an accidental or undetermined manner of death.

Requests for records were sent to the medical examiner's office, medical facilities, behavioral health agencies, law enforcement, and emergency medical services. All records received by the OFR Team were reviewed to determine if additional records were needed to analyze deaths and contributing factors.

Using these records, a case narrative was created for each case to be reviewed by the OFR team. Case reviews examined a person's demographics, psychosocial history, treatment history, medical records, crisis system encounters, and other prominent risk factors associated with drug overdoses.

From this, the team aims to identify missing gaps and opportunities for intervention. The outcome is to strengthen overdose prevention strategies, improve system-level operations, inform local service providers, public policy, and ultimately to reduce the number of overdose deaths in Mohave County.

There were five, two-hour reviews conducted by the Mohave County OFR Team. Case selection was limited to no more than seven cases per review meeting. A total of 27 cases were reviewed.

## Data Sources


Mohave County Department of Public Health collects data from a variety of sources including medical examiner data, toxicology reports, Controlled Substance Prescription Monitoring Program (CSPMP), pharmacy data, hospital records, Arizona Health Care Cost Containment System (AHCCCS), first responders, law enforcement, and other health care agencies.

# DATA OVERVIEW


## Demographics

As shown in Figures 1-3, most of the 27 decedents were male (85%). Most of the decedents were in the 25-34 age range (34%) and were white (74%). Additionally, only one of the decedents in our sample had prior military service (4%).


**Figure 1. Assigned Sex at Birth**





**Figure 2. Age**



**Figure 3. Race/Ethnicity**



As shown in Figures 4 and 5, the majority of decedents in our sample either graduated High School or received a GED certification (59%). Thirty-seven percent of decedents had some high school education but did not obtain a diploma or GED. Furthermore, 52% of the decedents were employed compared to 37% that were unemployed.




## Circumstances


Most of the fatal overdoses occurred at the decedent's place of residence (56%, Fig. 6). In 44% of the cases, the overdoses occurred in another residential location such as the home of a friend or family member or a public place, outdoor, hospital, or an automobile. More fatal overdoses occurred when a decedent was alone compared to with other adults present or nearby at the time of death (85% vs. 15%, Fig. 7). Naloxone was administered to 26% of decedents (Fig. 8) either by EMS or a bystander. Naloxone was administered to resuscitate from suspected overdose. Fifty-two percent of overdose deaths were found to be beyond resuscitative measures.




**Figure 6. Overdose Deaths by Injury Location**



**Figure 7. Presence Of A Witness**




**Figure 8. Use Of Naloxone**




In 78% of the cases, a combination of two or more substances were listed as a cause of death (Fig. 9). Forty-one percent of cases involved a combination of opioids and psychostimulants most frequently methamphetamine and fentanyl (Fig. 9). Methamphetamine was the most frequently found substance in the overdose deaths reviewed, present in 70% of cases, followed closely by fentanyl (67% Fig. 10). There was an increase in the presence of methamphetamine, while other substances such as heroin and other opioids decreased in 2023 compared to 2022.



**Figure 9. Single Substance vs. Combined Drug**



**Figure 10. Substances Contributing to Death**




## Contributing Factors & Commonalities


Multiple commonalities were noted to be shared among the sample of decedents. Commonalities are defined as shared attributes, circumstances, or life events shared by cases. They provide insight into risk factors and offer opportunities for intervention.

Many of the decedents had a mental health disorder (60%) or a chronic health condition (78%, Figure 11 & 12).-Depression (44%) and anxiety (44%) were the most prevalent mental health diagnoses. The most common health conditions were hypertension (30%) and chronic pain (30%). These numbers have the potential to be higher due to data limitations. Other attributes include (Table 1):

**Figure 11. Presence of a Mental Health Condition**



**Figure 12. Presence Of Chronic Health Conditions**




**Table 1. Other Attributes**


37%	Had a history of suicidal ideation and/or suicide attempts.
60%	Had a mental health condition.
78%	Had a chronic health condition.
33%	Had a previous non-fatal drug overdose.
30%	Recent incarceration release (past year).
41%	Had a history of incarceration at any point in their lifetime.
37%	Were homeless or had inadequate housing.

In 2023, we were able to obtain ACEs risk factors from eight out of 27 decedents (30%). Table 2 shows the percentage of ACEs risk factors, the highest being 50% had experienced physical abuse, emotional abuse, substance abuse within their household growing up, and/or parental separation or divorce. Additionally, Figure 13 shows the percentage of the decedents' age of first use of any substance, (at least 37% under the age of 18).

**Table 2. ACEs Risk Factors**



\*More than one risk factor may have been identified for each decedent.



## Controlled Substance Prescription Monitoring Program (CSPMP)


Of the 27 cases reviewed, almost half (48%) of the decedents had a history in the CSPMP. Five out of 13 of those with a CSPMP history had active prescriptions at the time of death, with only one of those having prescriptions that matched the drugs in the cause of death (Table 3).

**Table 3. History of Controlled Substance Prescriptions**

Characteristic	CSPMP, N (%)
History of prescription(s)	13 (48%)
Active prescription(s) at time of death	5 (19%)
Prescription(s) match drug(s) in COD	1 (4%)

## Prevention Recommendations

The Mohave County OFR team identified prevention recommendations for each of the 28 fatalities reviewed. Table 4 summarizes our recommendations for the sample along with the standardized recommendations.



<b>Prevalence of Prevention Recommendations by Base</b>		
<b>Healthcare</b>	<b>#</b>	<b>%</b>
Enhance capacity and coverage of mental health services (e.g., long-term case management).	12	44%
Enhance capacity and coverage of substance use disorder treatment (e.g., long-term treatment).	15	56%
Improve access to primary care services that include substance use disorder screening, harm reduction education, Medication Assisted Treatment, and referral to treatment.	9	33%
Establish protocols for emergency department and inpatient hospital discharge instructions to include naloxone distribution, connection to peer recovery specialists, case management, and treatment services.	6	22%
Increase patient-specific education of overdose risk associated with age, comorbid conditions, prescription medications, and the use of other substances.	16	59%
Improve care coordination by connecting to social support resources such as housing, employment insurance, transportation, etc.	5	19%
<b>Criminal Justice</b>		
Decrease stigma related to mental health and substance use disorders for incarcerated patients in an effort to increase the number of patients utilizing the available services.	4	15%
Improve care coordination post-release by performing a warm hand-off to substance use disorder treatment, medical care, and mental health services as well as providing social support resources such as housing, employment, insurance, transportation, etc.	5	19%
Increase naloxone distribution from correctional setting post-release.	4	15%
Increase training on evidence-based overdose prevention for corrections health and custody staff.	1	4%
Enhance capacity of drug courts to divert a greater number of individuals with a substance use disorder from incarceration.	2	7%
<b>Crisis Response</b>		
Improve care coordination by connecting to mental health and substance use services (e.g., transport/referral to treatment).	2	7%
Enhance universal training on evidence-based overdose prevention for first responders (EMS, Fire, Law Enforcement, etc.).	2	7%
Increase funding for crisis response teams that offer alternatives to incarceration.	0	0%
<b>Community/Public Health</b>		
Improve access to social support resources such as housing, employment, insurance, transportation, etc.	10	37%
Increase education and awareness of overdose prevention and harm reduction strategies (e.g., naloxone, fentanyl test strips, HIV/HEP C testing).	15	56%
Improve support for individuals with past childhood or adult trauma (e.g., domestic violence, sexual assault, loss of a loved one).	4	15%
Improve support for families who have a loved one with a substance use disorder (e.g., community events, connection to support groups, written materials).	18	67%
Develop social work/case management resources for children directly involved in overdose deaths to identify and treat behavioral health illness related to the death (e.g., PTSD from a parent's overdose death).	3	11%
<b>Fatality Review (if information missing or inadequate)</b>		
Promote best practice for drug-related death investigations.	0	0%
Promote the utilization of the ADHS State Lab comprehensive toxicology testing.	0	0%
Recruit new members to offer valuable insight at the community, system, or policy level.	1	4%

## Specific Prevention Recommendations for Mohave County

The top five recommendations (see below) were identified and shared with a variety of community partners to improve and enhance services in Mohave County.

### Healthcare

1. Enhance capacity and coverage of mental health services (e.g., long-term case management). (44%)
2. Enhance capacity and coverage of substance use disorder treatment (e.g., long-term treatment). (56%)
3. Increase patient-specific education of overdose risk associated with age, comorbid conditions, prescription medications, and the use of other substances. (56%)

### Community/Public Health

1. Increase education and awareness of overdose prevention and harm reduction strategies (e.g., naloxone, fentanyl test strips, HIV/HEP C testing). (56%)
2. Improve support for families who have a loved one with a substance use disorder (e.g., community events, connection to support groups, written materials). (67%)

## Conclusion

With the help of local stakeholders, Mohave County completed its fourth year of case reviews and created the OFR Annual Report. Deaths that occurred in 2023 were reviewed for this report.

The OFR's key recommendations are to enhance capacity and coverage of mental health services (e.g., long-term case management); enhance capacity and coverage of substance use disorder treatment (e.g., long-term treatment); increase patient-specific education of overdose risk associated with age, comorbid conditions, prescription medications, and the use of other substances; increase education and awareness of overdose prevention and harm reduction strategies (e.g., naloxone, fentanyl test strips, HIV/HEP C testing); and improve support for families who have a loved one with a substance use disorder (e.g., community events, connection to support groups, written materials).

The OFR findings and recommendations confirm the importance of collaboration and integrated care that identifies and takes advantage of multiple support services that engage individuals at risk of overdose.

The OFR team is committed to the ongoing analysis that identifies both gaps and prevention opportunities. The Mohave County Department of Public Health and the OFR team is committed to playing a positive role in reducing overdose deaths and encouraging healing among individuals who suffer from substance use disorder and their families by promoting evidence-based overdose prevention strategies identified by the Centers for Disease Control and Prevention (CDC).